

POLYGLYCOLIC ACID (PGA)

DESCRIPTION

SURCRYL is a synthetic, absorbable, sterile surgical suture composed of braided Polyglycolic Acid.

SURCRYL is colored violet with the dyestuff D&C Violet N°2. SURCRYL fulfills all the requirements of the European Pharmacopoeia and of the United States Pharmacopoeia.

INDICATIONS

SURCRYL suture is indicated for use in general soft tissue approximation and/or ligation, including use in ophthalmic procedures, but not for use in cardiovascular and neurological tissues.

ACTIONS

SURCRYL elicits a minimal acute inflammatory reaction in tissues, which is followed by gradual encapsulation of the suture by fibrous connective tissue. Loss of tensile strength and eventual absorption of Polyglycolic Acid (PGA) synthetic absorbable sutures occurs progressively. Absorption begins as a loss of tensile strength followed by a loss of mass. Implantation studies in animals indicate that Polyglycolic Acid (PGA) retains approximately 70% of its original tensile strength at two weeks post implantation, 50% of its original tensile strength at third week post implantation, and 20% of its original tensile strength at four weeks post implantation. Absorption of Polyglycolic Acid (PGA) absorbable synthetic suture is essentially complete between 60 and 90 days.

CONTRAINDICATIONS

This suture, being absorbable, should not be used where extended approximation of tissue is required. The use of this suture may be inappropriate in elderly, malnourished, or debilitated patients, or in patients suffering from conditions which may delay wound healing.

Not for use in cardiovascular and neurological surgery.

WARNINGS

Users should be familiar with surgical procedures and techniques involving absorbable sutures before employing SURCRYL suture for wound closure, as a risk of wound dehiscence may vary with the site of application and the suture material used. Do not use if package is open or damaged or the expiration date has been exceeded. Discard open, unused suture.

Do not resterilize; resterilization may alter the physical properties of this suture. Users should exercise caution when handling surgical needles to avoid inadvertent needle sticks. Discard used needles in a "sharps" container.

Avoid storing product at elevated temperatures.

As with any foreign body, prolonged contact of any suture with salt solutions, such as those found in the urinary or biliary tracts may result in calculus formation. As an absorbable suture, SURCRYL suture may act transiently as a foreign body.

Acceptable surgical practice should be followed for the management of contaminated or infected wounds.

PRECAUTIONS

Skin sutures, which must remain in place longer than 7 days may cause localized irritation and should be snipped off or removed as indicated. In handling this or any other suture material, care should be taken to avoid damage from handling. Avoid crushing or crimping damage due to application of surgical instruments such as forceps or needle holders.

Adequate knot security requires the accepted surgical technique of flat, square ties, with additional throws as warranted by surgical circumstances and the experience of the surgeon.

Users should be familiar with surgical procedures and techniques involving absorbable suture before employing SURCRYL synthetic absorbable suture for wound closure, as risk of wound dehiscence may vary with the site of application and the suture material used.

ADVERSE REACTIONS

Adverse effects associated with the use of this device include wound dehiscence, failure to provide adequate wound support in closure of the sites where expansion, stretching, or distension occur, failure to provide adequate wound support in elderly, malnourished or debilitated patients or in patients suffering from conditions which may delay wound healing, infection, minimal acute inflammatory tissue reaction, localized irritation when skin sutures are left in place for greater than 7 days, suture extrusion and delayed absorption in tissue with poor blood supply, calculi formation in urinary and biliary tracts when prolonged contact with salt solutions such as urine and bile occurs, and transitory local irritation at the wound site.

HOW SUPPLIED



SURCRYL sutures are available sterile, dyed and attached to stainless steel needles of varying types and sizes in one dozen boxes, or non-needled.

SURCRYL sutures are available in various lengths in sizes: 6/0 to 2 (0.7 to 5.0 metric).

CAUTION

Federal (USA) law restricts this device to sale by or on the order of a physician or licensed practitioner.

SYMBOLS USED ON LABELING

	CE symbol and identification number of the notified body.
	Lot number (batch)
	"Use by" expiration date (year-month)
	Date of manufacture (year-month)
	Do not use if package is damaged
	Single use
	consult instruction for use on
	Do not resterilize
	Manufacturer
	Product code
	Store below 25 C
	Keep away from sunlight
	protect from humidity
	Double sterile barrier system
	Medical Device
	Package is sterile when not damaged or opened. EO sterilization method -Ethylene oxide.



SURGIMEDIC

104 - 106 A6 Industrial Zone
10th of Ramadan-Egypt
T: +20 (0) 12 2219 4006
F: +20 (0) 2 2419 8050
E: contact@surgimed.com



Qualitech International B.V.

Siriusdreef 17
2132WT Hoofddorp
The Netherlands
Tel: +31207038302
mahdy@qt-int.uk